

United States
Environmental Protection
Agency


Office of Emergency and
Remedial Response
Washington DC 20460

Office of Research and Development
Office of Health and Environmental
Assessment
Environmental Criteria and
Assessment Office
Cincinnati OH 45268

Superfund



HEALTH EFFECTS ASSESSMENT FOR BENZO(a)PYRENE



HW
EPAX
8605-
0067

EPA/540/1-86-022
September 1984

**HEALTH EFFECTS ASSESSMENT
FOR BENZO(A)PYRENE**

**U.S. Environmental Protection Agency
Office of Research and Development
Office of Health and Environmental Assessment
Environmental Criteria and Assessment Office
Cincinnati, OH 45268**

**U.S. Environmental Protection Agency
Office of Emergency and Remedial Response
Office of Solid Waste and Emergency Response
Washington, DC 20460**

**U.S. Environmental Protection Agency
Region V, Library
230 South Dearborn Street
Chicago, Illinois 60604**

DISCLAIMER

This report has been funded wholly or in part by the United States Environmental Protection Agency under Contract No. 68-03-3112 to Syracuse Research Corporation. It has been subject to the Agency's peer and administrative review, and it has been approved for publication as an EPA document. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

U.S. Environmental Protection Agency

PREFACE

This report summarizes and evaluates information relevant to a preliminary interim assessment of adverse health effects associated with benzo(a)-pyrene. All estimates of acceptable intakes and carcinogenic potency presented in this document should be considered as preliminary and reflect limited resources allocated to this project. Pertinent toxicologic and environmental data were located through on line literature searches of the Chemical Abstracts, TOXLINE, CANCERLINE and the CHEMFATE/DATALOG data bases. The basic literature searched supporting this document is current up to September, 1984. Secondary sources of the information have also been relied upon in the preparation of this report and represent large scale health assessment efforts that entail extensive peer and Agency review. The following Office of Health and Environmental Assessment (OHEA) sources have been extensively utilized:

U.S. EPA. 1980b. Ambient Water Quality Criteria for Polynuclear Aromatic Hydrocarbons. Environmental Criteria and Assessment Office, Cincinnati, OH. EPA 440/5-80-069. NTIS PB 81-117806.

U.S. EPA. 1981. Hazard Profile for PAH. Prepared by the Environmental Criteria and Assessment Office, Cincinnati, OH, OHEA for the Office of Solid Waste and Emergency Response, Washington, DC.

U.S. EPA. 1983a. Review of Toxicologic Data in Support of Evaluation for Carcinogenic Potential of: Benzo[a]pyrene. Prepared by the Carcinogen Assessment Group, OHEA, Washington, DC for the Office of Solid Waste and Emergency Response, Washington, DC.

U.S. EPA. 1983b. Reportable Quantity for Benzo[a]pyrene. Prepared by the Environmental Criteria and Assessment Office, Cincinnati, OH, OHEA for the Office of Solid Waste and Emergency Response, Washington, DC.

The intent in these assessments is to suggest acceptable exposure levels whenever sufficient data were available. Values were not derived or larger uncertainty factors were employed when the variable data was limited in scope tending to generate conservative (i.e. protective) estimates. Nevertheless, the interim values presented reflect the relative degree of hazard associated with exposure or risk to the chemical(s) addressed.

Whenever possible, two categories of values have been estimated for systemic toxicants (toxicants for which cancer is not the endpoint of concern). The first, the AIS or acceptable intake subchronic, is an estimate of an exposure level which would not be expected to cause adverse effects when exposure occurs during a limited time interval (i.e., for an interval which does not constitute a significant portion of the lifespan). This type of exposure estimate has not been extensively used, or rigorously defined, as previous risk assessment efforts have been primarily directed towards exposures from toxicants in ambient air or water where lifetime exposure is assumed. Animal data used for AIS estimates generally include exposures with durations of 30-90 days. Subchronic human data are rarely available. Reported exposures are usually from chronic occupational exposure situations or from reports of acute accidental exposure.

The AIC, acceptable intake chronic, is similar in concept to the ADI (acceptable daily intake). It is an estimate of an exposure level that would not be expected to cause adverse effects when exposure occurs for a significant portion of the lifespan [see U.S. EPA (1980a) for a discussion of this concept]. The AIC is route specific and estimates acceptable exposure for a given route with the implicit assumption that exposure by other routes is insignificant.

Composite scores (CS) for noncarcinogens have also been calculated where data permitted. These values are used for ranking reportable quantities; the methodology for their development is explained in U.S. EPA (1983c).

For compounds for which there is sufficient evidence of carcinogenicity, AIS and AIC values are not derived. For a discussion of risk assessment methodology for carcinogens refer to U.S. EPA (1980a). Since cancer is a process that is not characterized by a threshold, any exposure contributes an increment of risk. Consequently, derivation of AIS and AIC values would be inappropriate. For carcinogens, q1*s have been computed based on oral and inhalation data if available.

ABSTRACT

In order to place the risk assessment evaluation in proper context, the reader is referred to the preface of this document. The preface outlines limitations applicable to all documents of this series as well as the appropriate interpretation and use of the quantitative estimates presented.

Animal data indicate that benzo(a)pyrene is a potent carcinogen via inhalation, oral and dermal routes. Human data concerning exposure to benzo(a)pyrene and cancer are lacking; however, human data concerning increased cancer risk and exposure to PAH containing mixtures are convincing.

U.S. EPA (1980b) used the mouse data of Neal and Rigdon (1967) to compute a q_1^* of $11.53 \text{ (mg/kg/day)}^{-1}$. This risk assessment has been extensively peer reviewed.

The same methodology was used in the present document to compute a q_1^* of $6.11 \text{ (mg/kg/day)}^{-1}$ from inhalation exposure data in Golden Syrian hamsters.

ACKNOWLEDGEMENTS

The initial draft of this report was prepared by Syracuse Research Corporation under Contract No. 68-03-3112 for EPA's Environmental Criteria and Assessment Office, Cincinnati, OH. Dr. Christopher DeRosa and Karen Blackburn were the Technical Project Monitors and Helen Ball was the Project Officer. The final documents in this series were prepared for the Office of Emergency and Remedial Response, Washington, DC.

Scientists from the following U.S. EPA offices provided review comments for this document series:

Environmental Criteria and Assessment Office, Cincinnati, OH
Carcinogen Assessment Group
Office of Air Quality Planning and Standards
Office of Solid Waste
Office of Toxic Substances
Office of Drinking Water

Editorial review for the document series was provided by:

Judith Olsen and Erma Durden
Environmental Criteria and Assessment Office
Cincinnati, OH

Technical support services for the document series was provided by:

Bette Zwyer, Pat Daunt, Karen Mann and Jacky Bohanon
Environmental Criteria and Assessment Office
Cincinnati, OH

TABLE OF CONTENTS

	<u>Page</u>
1. ENVIRONMENTAL CHEMISTRY AND FATE.	1
2. ABSORPTION FACTORS IN HUMANS AND EXPERIMENTAL ANIMALS	3
2.1. ORAL	3
2.2. INHALATION	3
3. TOXICITY IN HUMANS AND EXPERIMENTAL ANIMALS	4
3.1. SUBCHRONIC	4
3.1.1. Oral.	4
3.1.2. Inhalation.	4
3.2. CHRONIC.	4
3.2.1. Oral.	4
3.2.2. Inhalation.	4
3.3. TERATOGENICITY AND OTHER REPRODUCTIVE EFFECTS.	4
3.3.1. Oral.	4
3.3.2. Inhalation.	5
3.4. TOXICANT INTERACTIONS.	5
4. CARCINOGENICITY	7
4.1. HUMAN DATA	7
4.1.1. Oral.	7
4.1.2. Inhalation.	7
4.2. BIOASSAYS.	8
4.2.1. Oral.	8
4.2.2. Inhalation.	8
4.3. OTHER RELEVANT DATA.	15
4.4. WEIGHT OF EVIDENCE	16
5. REGULATORY STANDARDS AND CRITERIA	17

TABLE OF CONTENTS (cont.)

	<u>Page</u>
6. RISK ASSESSMENT	19
6.1. ACCEPTABLE INTAKE SUBCHRONIC (AIS)	19
6.2. ACCEPTABLE INTAKE CHRONIC (AIC).	19
6.3. CARCINOGENIC POTENCY (q_1^*)	19
6.3.1. Oral.	19
6.3.2. Inhalation.	19
7. REFERENCES.	21
APPENDIX A: Summary Table for Benzo[a]pyrene	28
APPENDIX B: Cancer Data Sheet for Derivation of q_1^*	29
APPENDIX C: Cancer Data Sheet for Derivation of q_1^*	31

LIST OF TABLES

<u>No.</u>	<u>Title</u>	<u>Page</u>
1-1	Selected Physical and Chemical Properties and Half-lives for Benzo[a]pyrene.	2
4-1	Carcinogenicity of Benzo[a]pyrene Administered in the Diet to Male and Female CFW Mice at Levels of 1-250 ppm.	9
4-2	Carcinogenicity of Benzo[a]pyrene Administered in the Diet to Male and Female Swiss Mice at Levels of 250-1000 ppm . .	10
4-3	Carcinogenicity of Benzo[a]pyrene Administered in the Diet to Male and Female Swiss CFW Mice at a Level of 250 ppm . .	11
4-4	Carcinogenicity of Benzo[a]pyrene to Male Syrian Golden Hamsters by Inhalation.	12
4-5	Carcinogenicity of Benzo[a]pyrene in Syrian Hamsters Following Intratracheal Administration of 0.10-1.0 mg/week.	13
4-6	Carcinogenicity of Benzo[a]pyrene in Golden Syrian Hamsters Following Intratracheal Administration of 18.2-36.4 mg/animal	14

LIST OF ABBREVIATIONS

ADI	Acceptable daily intake
AIC	Acceptable intake chronic
AIS	Acceptable intake subchronic
bw	Body weight
CAS	Chemical Abstract Service
CS	Composite score
DNA	Deoxyribonucleic acid
GI	Gastrointestinal
PAH	Polycyclic aromatic hydrocarbon
SCE	Sister chromatid exchange
TLV	Threshold limit value
TWA	Time-weighted average

1. ENVIRONMENTAL CHEMISTRY AND FATE

The relevant physical and chemical properties and environmental fate of benzo[a]pyrene (CAS Registry No. 50-32-8) are given in Table 1-1.

The half-life for benzo[a]pyrene in the atmosphere has been estimated from the discussion in the NAS (1983) document regarding the chemical and its photochemical reaction and the estimation of its half-life due to physical removal processes as discussed by Cupitt (1980).

Pertinent data regarding the leachability of this compound in soils could not be located in the available literature. Considering the high octanol/water partition coefficient and low water solubility, the compound is expected to have very low mobility in soils, particularly in soils with high organic matter content.

TABLE 1-1

Selected Physical and Chemical Properties and Half-lives for Benzo[a]pyrene

Properties	Values	Reference
Chemical class:	Polycyclic aromatic hydrocarbons (PAH)	NA
Molecular weight:	252	Mabey et al., 1981
Vapor pressure:	5.6×10^{-9} mm Hg at 25°C	Mabey et al., 1981
Water solubility:	1.2 µg/kg at 25°C	Wise et al., 1981
Log octanol/water partition coefficient:	6.06	U.S. EPA, 1980b
Bioconcentration factor:	28,200 (estimated)	U.S. EPA, 1980b
Half-lives in:		
Air:	<1-6 days (estimated)	NAS, 1983; Cupitt, 1980
Water:	<1-8 hours in aquatic phase 5-10 years in sediment	Smith et al., 1978 Smith et al., 1978
Soil:	>14-16 months for complete degradation	U.S. EPA, 1981

NA = Not applicable

2. ABSORPTION FACTORS IN HUMANS AND EXPERIMENTAL ANIMALS

2.1. ORAL

Data regarding oral absorption of benzo[a]pyrene are limited; however, observed toxicity following oral administration indicates that benzo[a]pyrene is absorbed by this route (Smyth et al., 1962; U.S. EPA, 1980b, 1981; Santodonato et al., 1981).

Absorption of orally administered lipophilic PAH can be hampered by the mucous layer lining the GI tract (Grimmer, 1983). Rats given benzo[a]pyrene in starch solution (100 mg) by gavage or in the diet (250 mg) absorbed ~50% of the administered compound (Chang, 1943).

Regardless of the type of solvent used, benzo[a]pyrene readily penetrates the forestomach epithelium of mice. In the glandular stomach, however, the type of solvent used plays a decisive role in the absorption of benzo[a]pyrene (Ekwall et al., 1951; Setälä, 1954). Hydrophilic solvents enhance the absorption of benzo[a]pyrene from the glandular stomach as compared to lipophilic solvents.

2.2. INHALATION

Data regarding the pulmonary absorption of benzo[a]pyrene are limited; however, observed toxicity after inhalation exposure indicates that benzo[a]pyrene is readily absorbed through the lungs (Kotin et al., 1969; Vainio et al., 1976). As a class, PAHs are highly lipid soluble and capable of passing across epithelial membranes (U.S. EPA, 1980b).

3. TOXICITY IN HUMANS AND EXPERIMENTAL ANIMALS

3.1. SUBCHRONIC

3.1.1. Oral. Differences in genetic constitutions appear to influence the subchronic oral toxicity of benzo[a]pyrene in mice. Specifically, the Ah locus, which determines the inducibility of aryl hydrocarbon hydroxylase, plays a major role in determining the oral toxicity of benzo[a]pyrene, presumably by influencing the pathways of biotransformation. Robinson et al. (1975) administered benzo[a]pyrene in the diet at a level of 120 mg/kg bw to nonresponsive (poorly inducible) AKR/N mice (Ah^d/Ah^d type) and to responsive (markedly inducible) mice (Ah^b/Ah^b type). Nonresponsive mice developed aplastic anemia and died within 4 weeks, whereas responsive mice remained healthy for at least 6 months.

3.1.2. Inhalation. Pertinent data regarding the non-tumor-related subchronic toxicity of benzo[a]pyrene administered by inhalation could not be located in the available literature.

3.2. CHRONIC

3.2.1. Oral. The only available chronic bioassays for oral exposure of benzo[a]pyrene are investigations of carcinogenicity (U.S. EPA, 1980b). The lack of appropriate protocols (i.e., non-tumor pathology) and detailed reporting of symptoms render these carcinogenicity bioassays inadequate for use in evaluating non-carcinogenic endpoints.

3.2.2. Inhalation. There are no reports available concerning the non-tumor-related chronic toxicity of benzo[a]pyrene administered by inhalation.

3.3. TERATOGENICITY AND OTHER REPRODUCTIVE EFFECTS

3.3.1. Oral. Rigdon and Rennels (1964) fed female rats a diet containing benzo[a]pyrene at a level of 1000 mg/kg (equivalent to 50 mg/kg/day) for up to 3.5 months. Of seven pregnant treated animals, only one dam carried viable fetuses to term, delivering four pups on the 23rd day of pregnancy.

Two of the four pups were stillborn, one of which was grossly malformed (not necessarily treatment-related). A third pup was killed for observational purposes, and evidently the fourth pup died of starvation 3 days after birth because the dam did not appear to be lactating. The authors were not certain whether this absence of lactation was treatment-related.

At autopsy, four dead fetuses were found in the right uterine horn of a second dam. Signs of toxicity (body weight changes or histopathological changes) were not observed in the treated dams.

In a teratogenicity and reproduction study, Rigdon and Neal (1965) fed male and female mice diets containing benzo[a]pyrene at a level of 0, 250, 500 or 1000 mg/kg over various time spans during mating, gestation and lactation. No apparent reproductive, teratogenic, embryotoxic or fetotoxic effects were observed in the experimental animals.

Mackenzie and Angevine (1981) administered benzo[a]pyrene orally at a level of 10 mg/kg bw to CD-1 mice during pregnancy. There was no effect on fetal body weights, but there was a marked and specific reduction of gonadal weight, and reduced fertility and reproductive capacity were observed among the offspring. At a level of 40 mg/kg/day, almost complete sterility was observed in both sexes of offspring.

3.3.2. Inhalation. Pertinent data regarding the teratogenic effects resulting from inhalation exposure to benzo[a]pyrene could not be located in the available literature.

3.4. TOXICANT INTERACTIONS

U.S. EPA (1980b) has described extensively the synergistic and antagonistic interactions among different PAHs and between PAH and non-PAH chemicals. Briefly, metabolism of PAH by the microsomal mixed function oxidase

enzyme system yields several types of reactive and potentially carcinogenic intermediates. Chemicals that induce or inhibit this enzyme system alter the patterns of PAH metabolism and, hence, alter their toxic and carcinogenic properties.

4. CARCINOGENICITY

The carcinogenicity of benzo[a]pyrene has been tested extensively by applying it to the skin of mice, and only infrequently by other routes of administration. The studies discussed below have been summarized in U.S. EPA (1983a); however, more complete reviews of the carcinogenicity bioassays of benzo[a]pyrene are provided by IARC (1973, 1983), U.S. EPA (1980b, 1981, 1983b) and Santodonato et al. (1981).

4.1. HUMAN DATA

Few case reports are available regarding the direct carcinogenic effects of benzo[a]pyrene on humans. Cottini and Mazzone (1939) applied a 1% solution of benzo[a]pyrene in benzene to small areas of the exposed and unexposed skin of 26 patients. Up to 120 daily applications were applied over a 4-month period, within which time, regressive verrucae developed in each of the 26 patients. Although reversible and apparently benign, these changes were thought to represent early stages of neoplastic proliferation. Similar cases of epidermal changes were reported by Rhoads et al. (1954) and Klar (1938) to have occurred in men accidentally exposed to benzo[a]pyrene. Numerous epidemiologic studies of human populations (primarily worker groups) have shown a clear association between exposure to PAH-containing materials (e.g., soots, tars, oils) and increased cancer risk (Santodonato et al., 1981; IARC, 1973, 1983; U.S. EPA, 1981).

4.1.1. Oral. Pertinent data regarding the carcinogenicity of benzo[a]pyrene to humans following oral exposure could not be located in the available literature.

4.1.2. Inhalation. Pertinent data regarding the carcinogenicity of benzo[a]pyrene to humans following inhalation exposure could not be located in the available literature.

4.2. BIOASSAYS

4.2.1. Oral. Benzo[a]pyrene was administered in various concentrations in the diets of mice to test its carcinogenicity (Neal and Rigdon, 1967; Rigdon and Neal, 1966, 1969). These studies are summarized in Tables 4-1, 4-2 and 4-3. A dose-response relationship was noted for the incidence of stomach tumors (papillomas and carcinomas) in male and female CFW-Swiss mice treated with 1-250 ppm benzo[a]pyrene for up to 197 days (Neal and Rigdon, 1967). Stomach tumors were reported in animals treated with 20, 40, 45, 50, 100 and 250 ppm benzo[a]pyrene (5/23, 1/40, 4/40, 24/34, 19/23 and 66/73, respectively), while control animals (0/289) and those treated with 1, 10 and 30 ppm benzo[a]pyrene (0/25, 0/24 and 0/37, respectively) did not have stomach tumors. In addition to the increase in stomach tumors, increased incidences of lung adenoma and leukemia were noted in mice treated with 250 and 1000 ppm benzo[a]pyrene (Rigdon and Neal, 1966, 1969).

4.2.2. Inhalation. Thyssen et al. (1981) exposed groups of 24 hamsters by inhalation to benzo[a]pyrene at levels of 2.2, 9.5 or 45 mg/m³ for 4.5 hours/day for 10 weeks and 3 hours/day 7 days/week thereafter, for up to 675 days (Table 4-4). No treatment-related tumors were observed in animals exposed to 2.2 mg/m³. Animals exposed to 9.5 mg/m³, however, developed tumors of the nasal cavity (12%), larynx (31%), trachea (4%) and pharynx (23%). Hamsters exposed to 44.8 mg/m³ benzo[a]pyrene developed tumors of the respiratory tract (13/25) and upper digestive tract (14/25). No tumors of these types were seen in control animals (Thyssen et al., 1981).

Intratracheal administration of benzo[a]pyrene resulted in an increased incidence of respiratory tract neoplasms in both sexes of Syrian hamsters (Ketkar et al., 1978; Feron and Kruysse, 1978) (Tables 4-5 and 4-6). A dose-related response was reported for hamsters treated with 18.2 and 36.4 mg/animal (total dose) for 52 weeks followed by a 29-week latency period.

TABLE 4-1

Carcinogenicity of Benzo[a]pyrene Administered in the Diet to Male and Female CFM Mice at Levels of 1-250 ppm*

Dose	Duration of Treatment (days)	Duration of Study (days)	Purity of Compound	Vehicle or Physical State	Target Organ	Tumor Type	Tumor Incidence
1 ppm (0.48 mg total dose)	110	140	NR	diet	stomach	papillomas/carcinomas	0/25
10 ppm (4.48 mg total dose)	110	140	NR	diet	stomach	papillomas/carcinomas	0/24
20 ppm (8.88 mg total dose)	110	226	NR	diet	stomach	papillomas/carcinomas	5/23
30 ppm (13.32 mg total dose)	110	143-177	NR	diet	stomach	papillomas/carcinomas	0/37
40 ppm (17.76 mg total dose)	110	143-211	NR	diet	stomach	papillomas/carcinomas	1/40
45 ppm (19.8 mg total dose)	110	141-183	NR	diet	stomach	papillomas/carcinomas	4/40
50 ppm (21.4-29.4 mg total dose)	107-197	124-219	NR	diet	stomach	papillomas/carcinomas	24/34
100 ppm (39.2-48.8 mg total dose)	98-122	118-146	NR	diet	stomach	papillomas/carcinomas	19/23
250 ppm (70-165 mg total dose)	70-165	88-185	NR	diet	stomach	papillomas/carcinomas	66/73
0.0 ppm	NA	70-300	NA	basal diet only	stomach	papillomas/carcinomas	0/289

*Source: Neal and Rigdon, 1967

NA - Not applicable; NR - Not reported

TABLE 4-2

Carcinogenicity of Benzo[a]pyrene Administered in the Diet to Male and Female Swiss Mice at Levels of 250-1000 ppm*

Dose	Duration of Treatment (days)	Duration of Study (days)	Purity of Compound	Vehicle or Physical State	Target Organ	Tumor Type	Tumor Incidence
1000 ppm (1 mg/g food)	73-83	73-83	NR	diet	stomach lung	papilloma/carcinoma adenoma	5/9 7/9
1000 ppm (1 mg/g food)	127-187	127-187	NR	diet	stomach lung	papilloma/carcinoma adenoma	13/13 3/13
250 ppm (0.25 mg/g food)	72-99	72-99	NR	diet	stomach lung	papilloma/carcinoma adenoma	12/52 26/52
250 ppm (0.25 mg/g food)	147-196	147-196	NR	diet	stomach lung	papilloma/carcinoma adenoma	9/13 10/13
0.0 ppm	NA	111-120	NA	diet only	stomach lung	papilloma/carcinoma adenoma	2/108 25/108

*Source: Rigdon and Neal, 1966

NA = Not applicable; NR = Not reported

TABLE 4-3

Carcinogenicity of Benzo[a]pyrene Administered in the Diet to Male and Female Swiss CFW Mice at a Level of 250 ppm^a

Dose	Duration of Treatment	Duration of Study	Purity of Compound	Vehicle or Physical State	Target Organ	Tumor Type	Tumor Incidence
250 ppm (0.25 mg/g food)	80-140 days	80-140 days	NR	diet	stomach	papilloma/carcinoma	69/108
					lung	adenoma	52/108
					hematopoietic system	leukemia	40/108
0.0 ppm	NA	62-300 days	NA	diet only	stomach	papilloma/carcinoma	2/175 ^b
					lung	adenoma	33/151
					hematopoietic system	leukemia	0/175 ^b

^aSource: Rigdon and Neal, 1969^bIncidence of tumors in a control group reported previously by Rigdon and Neal (1966).

NA - Not applicable; NR - Not reported

TABLE 4-4

Carcinogenicity of Benzo[a]pyrene to Male Syrian Golden Hamsters by Inhalation^{a,b}

Dose	Duration of Treatment (weeks)	Duration of Study (weeks)	Purity of Compound	Vehicle or Physical State	Target Organ	Tumor Type ^c	Tumor Incidence
2.2 mg/m ^a (29 mg total dose)	95.2	95.2	NR	NaCl vapor in air	respiratory tract upper digestive tract	tumors tumors	0/27 0/27
9.5 mg/m ^a (127 mg total dose)	96.4	96.4	NR	NaCl vapor in air	respiratory tract upper digestive tract	tumors tumors	9/26 ^d 7/26 ^d
46.5 mg/m ^a (383 mg total dose)	59.5	59.5	NR	NaCl vapor in air	respiratory tract upper digestive tract	tumors tumors	13/25 ^e 14/25 ^e
0.0 mg/m ^a	NA	96.4	NA	NaCl vapor only	respiratory tract upper digestive tract	tumors tumors	0/27 0/27

^aSource: Thyssen et al., 1981^bExposure was for 4.5 hours/day for the first 10 weeks, 3 hours/day thereafter for 7 days/week.^cTumors were papillomas, papillary polyps, and squamous cell carcinomas.^d3 nasal cavity, 8 laryngeal, 1 tracheal, 6 pharyngeal and 1 forestomach tumors^e1 nasal cavity, 13 laryngeal, 3 tracheal, 14 pharyngeal, 2 esophageal and 1 forestomach tumor

NA = Not applicable; NR = Not reported

TABLE 4-5

Carcinogenicity of Benzo[a]pyrene in Syrian Hamsters Following Intratracheal Administration of 0.10-1.0 mg/week^a

Sex	Dose (mg/week)	Duration of Treatment ^b (weeks)	Duration of Study ^b (weeks)	Purity of Compound	Vehicle or Physical State	Target Organ	Tumor Type ^c	Tumor Incidence
M	0.10	40	40	97%	bovine albumin	respiratory tract	various neoplasms	5/26
F	0.10	34	34	97%	bovine albumin	respiratory tract	various neoplasms	12/30
M	0.33	24	24	97%	bovine albumin	respiratory tract	various neoplasms	7/29
F	0.33	28	28	97%	bovine albumin	respiratory tract	various neoplasms	10/28
M	1.0	10	10	97%	bovine albumin	respiratory tract	various neoplasms	6/27
F	1.0	15	15	97%	bovine albumin	respiratory tract	various neoplasms	6/30
M	0.0	41	41	NA	bovine albumin only	respiratory tract	various neoplasms	0/29
F	0.0	35	35	NA	bovine albumin only	respiratory tract	various neoplasms	0/30

^aSource: Ketkar et al., 1978^bMean survival time^cCarcinomas, adenomas, adenocarcinomas and papillomas were reported.

NA = Not applicable

TABLE 4-6

Carcinogenicity of Benzo[a]pyrene in Golden Syrian Hamsters Following Intratracheal Administration of 18.2-36.4 mg/animal^a

Sex	Dose	Duration of Treatment (weeks)	Duration of Study (weeks)	Purity of Compound	Vehicle or Physical State	Target Organ	Tumor Type ^b	Tumor Incidence
M	18.2 mg/hamster total dose	52 (1 dose/week)	81	>99%	0.9% NaCl	respiratory tract	various	4/29
M	36.4 mg/hamster total dose	52 (1 dose/week)	81	>99%	0.9% NaCl	respiratory tract	various	19/30
M	0.0 mg/hamster total dose	52 (1 dose/week)	81	>99%	saline vehicle only	respiratory tract	various	0/30 ^c
F	18.2 mg/hamster total dose	52 (1 dose/week)	81	>99%	0.9% NaCl	respiratory tract	various	3/27
F	36.4 mg/hamster total dose	52 (1 dose/week)	81	>99%	0.9% NaCl	respiratory tract	various	7/24
F	0.0 mg/hamster total dose	52 (1 dose/week)	81	>99%	saline vehicle only	respiratory tract	various	0/28 ^c

^aSource: Feron and Kruysse, 1978^bPapillomas and carcinomas of the trachea and pulmonary adenomas were most prevalent.^cCombined tumor incidence of untreated and vehicle controls

The incidence of tracheal papillomas and carcinomas, collectively, with lung adenomas was 4/29 and 3/27 for low dose males and females, respectively, and 19/30 and 7/24 for high dose males and females, respectively (Feron and Kruyse, 1978). Ketkar et al. (1978) reported a high dose-related mortality in hamsters treated at dose levels lower than those used by Feron and Kruyse (1978). Mean survival times ranged from 40 weeks for male hamsters treated with 0.1 mg benzo[a]pyrene/week to 10 weeks for males treated with 1.0 mg benzo[a]pyrene/week. An increase in the incidence of respiratory tract carcinoma, adenoma and papilloma in both sexes of hamsters in the treatment groups was reported, but a definite dose-related response was not evident (Ketkar et al., 1978).

4.3. OTHER RELEVANT DATA

The mutagenicity of benzo[a]pyrene has been summarized by IARC (1982), U.S. EPA (1980b, 1981), Santodonato et al. (1981), deSerres and Ashby (1981) and Hollstein and McCann (1979). The reader is referred to these reviews for further information.

Benzo[a]pyrene is an indirect acting carcinogen, undergoing metabolism to a reactive electrophile capable of covalently binding to DNA (IARC, 1982; Lutz, 1979). Benzo[a]pyrene has been used extensively as a model carcinogen and as a positive control in a variety of short-term tests, yielding positive results in assays for bacterial DNA repair, bacteriophage induction and bacterial mutation, mutation in Drosophila melanogaster; DNA binding, DNA repair, SCE, chromosomal aberration, point mutation and transformation in mammalian cells in culture; and in tests in mammals in vivo, including DNA binding, SCE, chromosomal aberration, sperm abnormality and the specific locus (spot) test (IARC, 1982; deSerres and Ashby, 1981; Hollstein and McCann, 1979).

4.4. WEIGHT OF EVIDENCE

Benzo[a]pyrene is both a local and a systemic carcinogen, producing tumors in rats, mice, hamsters, guinea pigs, rabbits and monkeys following oral, inhalation or dermal exposure. Benzo[a]pyrene is an initiator of skin carcinogenesis in mice and also produces tumors following single doses or prenatal exposure. Benzo[a]pyrene has been used extensively as a model carcinogen and as a positive control in a variety of short-term tests. IARC (1982) reported that there is sufficient evidence that benzo[a]pyrene is an animal carcinogen and limited evidence that it is a human carcinogen. Applying the classification criteria for weight of evidence proposed by the Carcinogen Assessment Group of the U.S. EPA (Federal Register, 1984), benzo[a]pyrene is most appropriately designated a Group B2 chemical.

5. REGULATORY STANDARDS AND CRITERIA

Exposure criteria and TLVs have been developed for PAH as a class, as well as for several individual PAHs. The Occupational Safety and Health Administration has set an 8-hour TWA concentration limit of 0.2 mg/m³ for the benzene-soluble fraction of coal tar pitch volatiles (anthracene, benzo[a]pyrene, phenanthrene, acridine, chrysene, pyrene) (Code of Federal Regulations, 1981). NIOSH (1977) recommends a concentration limit for coal tar, coal tar pitch, creosote and mixtures of these substances at 0.1 mg/m³ of the cyclohexane-extractable fraction of the sample, determined as a 10-hour TWA. NIOSH (1977) concluded that these specific coal tar products, as well as coke oven emissions, are carcinogenic and can increase the risk of lung and skin cancer in workers. NIOSH (1977) also recommends a ceiling limit for exposure to asphalt fumes of 5 mg airborne particulates/m³ of air.

Environmental quality criteria for PAH have been recommended for ambient water, which specify concentration limits intended to protect humans against adverse health effects. The U.S. EPA (1980b) has recommended a concentration limit of 28 ng/l for the sum of all carcinogenic PAHs in ambient water. This value is based on a mathematical extrapolation of the results from studies with mice treated orally with benzo[a]pyrene, and acknowledges the conservative assumption that all carcinogenic PAHs are equal in potency to benzo[a]pyrene. Daily consumption of water containing 28 ng/l of carcinogenic PAH over an entire lifetime is estimated on the basis of the animal bioassay data to keep the lifetime risk of cancer development below 1 chance in 100,000.

The U.S. EPA has not recommended an ambient water quality criterion for noncarcinogenic PAH as a class. The U.S. EPA (1980b) acknowledged that data suitable for quantitative risk assessment of noncarcinogenic PAH are essentially nonexistent.

6. RISK ASSESSMENT

6.1. ACCEPTABLE INTAKE SUBCHRONIC (AIS)

Benzo[a]pyrene is a known carcinogen for which data are sufficient for computing a q_1^* . It is, therefore, inappropriate to calculate an oral or inhalation AIS for benzo[a]pyrene.

6.2. ACCEPTABLE INTAKE CHRONIC (AIC)

Benzo[a]pyrene is a known carcinogen for which data are sufficient for computing a q_1^* . It is, therefore, inappropriate to calculate an oral or inhalation AIC for benzo[a]pyrene.

6.3. CARCINOGENIC POTENCY (q_1^*)

6.3.1. Oral. A carcinogenic potency factor, q_1^* , for humans can be derived from the study of Neal and Rigdon (1967) in which benzo[a]pyrene at dose levels of 1-250 ppm in the diet was administered to strain CFW mice for ~110 days. The incidences of stomach tumors (mostly squamous-cell papillomas, but some carcinomas) were 0/289 for controls, 0/25 at the 1 ppm (0.13 mg/kg/day) level, 0/24 at 10 ppm (1.3 mg/kg/day), 1/23 at 20 ppm (2.6 mg/kg/day), 0/37 at 30 ppm (3.9 mg/kg/day), 1/40 at 40 ppm (5.2 mg/kg/day), 4/40 at 45 ppm (5.85 mg/kg/day), 24/34 at 50 ppm (6.5 mg/kg/day), 19/23 at 100 ppm (13.0 mg/kg/day) and 66/73 at 250 ppm (32.5 mg/kg/day). U.S. EPA (1980b) used these incidences of stomach tumors to derive a q_1^* of 11.53 (mg/kg/day)⁻¹. The data used in derivation of this q_1^* are presented in Appendix C.

6.3.2. Inhalation. A carcinogenic potency factor, q_1^* , for humans can be derived from the study of Thyssen et al. (1981) in which Golden Syrian hamsters were exposed to benzo[a]pyrene by inhalation at levels of 0, 2.2, 9.5 or 46.5 mg/m³ for 10-96.4 weeks. The incidences of respiratory tumors were 0/27 for controls, 0/27 for the low-dose group, 9/26 for the mid-dose

group and 13/25 for the high-dose group. The highest exposure level did have an adverse impact on survival time. Animals exposed to an average concentration of 46.5 mg benzo[a]pyrene/m³ air had an average survival time of 59.5 weeks, as compared with controls which survived for an average of 96.4 weeks. Due to early mortality in the highest dose group, these data were excluded from the q_1^* derivation. Based on the respiratory tumor response of male hamsters and using the linearized multistage model adopted by the U.S. EPA (U.S. EPA, 1980a), a carcinogenic potency factor (q_1^*) of $\checkmark 6.11 \text{ (mg/kg/day)}^{-1}$ can be derived for humans. The animal q_1^* was derived using transformed doses based on an inhalation rate of hamsters of 0.037 m³/day and assuming 100% absorption. The conversion factor used in the calculation of the human q_1^* from the animal data was the cube root of the body weight ratio, empirically accepted to adjust for species to species extrapolation. Complete data for derivation of the q_1^* are presented in Appendix B.

7. REFERENCES

Chang, L.H. 1943. The fecal excretion of polycyclic hydrocarbons following their administration to the rat. J. Biol. Chem. 151: 93. (Cited in Grimmer, 1983)

Code of Federal Regulations. 1981. OSHA Safety and Health Standards. 29 CFR 1910.1000.

Cottini, G.B. and G.B. Mazzone. 1939. The effects of 3,4-benzpyrene on human skin. Am. J. Cancer. 37: 186. (Cited in U.S. EPA, 1983a)

Cupitt, L. 1980. Fate of Toxic and Hazardous Materials in the Air Environment. Environmental Sciences Research Laboratory, ORD, U.S. EPA, Research Triangle Park, NC. EPA 600/3-80-084. NTIS PB 80-221948.

deSerres, F. and J. Ashby, Ed. 1981. Evaluation of Short-Term Tests for Carcinogens. Report of the International Collaborative Program. Elsevier/North-Holland Biomedical Press, New York. p. 180, 190. (Cited in IARC, 1982)

Ekwall, P., P. Ermala, K. Setala and L. Sjoblom. 1951. Gastric absorption of 3,4-benzpyrene. II. The significance of the solvent for the penetration of 3,4-benzpyrene into the stomach wall. Cancer Res. 11: 758. (Cited in Grimmer, 1983)

Federal Register. 1984. Environmental Protection Agency. Proposed guidelines for carcinogenic risk assessment. 49 FR 46294-46299.

Feron, V.J. and A. Kruysse. 1978. Effects of exposure to furfural vapour in hamsters simultaneously treated with benzo[a]pyrene or diethylnitrosamine. Toxicology. 11: 127-144. (Cited in U.S. EPA, 1983a)

Grimmer, G. 1983. Environmental Carcinogens: Polycyclic Aromatic Hydrocarbons. Chemistry, Occurrence, Biochemistry, Carcinogenicity. CRC Press, Inc., Boca Raton, FL. p. 27-60.

Hollstein, M. and J. McCann. 1979. Short-term tests for carcinogens and mutagens. Mutat. Res. 65: 133-226. (Cited in IARC, 1982)

IARC (International Agency for Research on Cancer). 1973. Certain polycyclic aromatic hydrocarbons and heterocyclic compounds. In: IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man. WHO, IARC, Lyon, France. Vol. 3.

IARC (International Agency for Research on Cancer). 1982. Chemicals, Industrial processes and Industries Associated with Cancer in Humans. In: IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. WHO, IARC, Lyon, France. Suppl. 4.

IARC (International Agency for Research on Cancer). 1983. Polynuclear Aromatic Compounds, Part 1, Chemical, Environmental and Experimental Data. In: IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. WHO, IARC, Lyon, France. Vol. 32.

Ketkar, M., G. Resnik, P. Schneider and U. Mohr. 1978. Investigations on the carcinogenic burden by air pollution in man. Intratracheal instillation studies with benzo[a]pyrene in bovine serum albumin in Syrian hamsters. Cancer Lett. 4(4): 235-239. (Cited in U.S. EPA, 1983a)

Klar, E. 1938. Über die entstehung eines epithelioms beim menschen nach experimentellen arbeiten mit benzpyren. Klin. Wschr. 17: 1279. (Ger.) (Cited in IARC, 1973; U.S. EPA, 1983a)

Kotin, P., H.L. Falk and R. Busser. 1969. Distribution, retention, and elimination of C¹⁴-3,4-benzpyrene after administration to mice and rats. J. Natl. Cancer Inst. 23: 541-555. (Cited in U.S. EPA, 1980b)

Lutz, W.K. 1979. In vivo covalent binding of organic chemicals to DNA as a quantitative indication in the process of chemical carcinogenesis. Mutat. Res. 65: 289-356. (Cited in IARC, 1982)

Mabey, W.R., J.H. Smith, R-T. Podoll, et al. 1981. Aquatic Fate Process Data for Organic Priority Pollutants. Monitoring and Data Support Division, Office of Water Regulations and Standards, Washington, DC. EPA 440/4-81-014.

MacKenzie, K.M. and D.M. Angevine. 1981. Infertility in mice exposed in utero to benzo[a]pyrene. Biol. Reprod. 24: 183-191. (Cited in IARC, 1983)

NAS (National Academy of Sciences). 1983. Polycyclic Aromatic Hydrocarbons: Evaluation of Sources and Effects. Committee on Pyrene and Selected Analogues. Board on Toxicology and Environmental Health Hazards, Commission on Life Sciences, National Research Council, Washington, DC.

Neal, J. and R.H. Rigdon. 1967. Gastric tumors in mice fed benzo[a]pyrene: A quantitative study. Tex. Rep. Biol. Med. 25: 553. (Cited in U.S. EPA, 1983a)

NIOSH (National Institute for Occupational Safety and Health). 1977. Criteria for a Recommended Standard...Occupational Exposure to Coal Tar Products. U.S. DHEW, PHS, CDC, Rockville, MD.

Rhoads, C.P., W.E. Smith, N.S. Cooper and R.D. Sullivan. 1954. Early changes in the skin of several species including man, after painting with carcinogenic materials. Proc. Am. Assoc. Cancer Res. 1: 40. (Cited in IARC, 1973; U.S. EPA, 1983a)

Rigdon, R.H. and J. Neal. 1965. Effects of feeding benzo(a)pyrene on fertility, embryos, and young mice. J. Natl. Cancer Inst. 34: 297-305. (Cited in U.S. EPA, 1983b)

Rigdon, R.H. and J. Neal. 1966. Gastric carcinomas and pulmonary adenomas in mice fed benzo[a]pyrene. Tex. Rep. Biol. Med. 24: 195. (Cited in U.S. EPA, 1983a)

Rigdon, R.H. and J. Neal. 1969. Relationship of leukemia to lung and stomach tumors in mice fed benzo[a]pyrene. Proc. Soc. Exp. Biol., NY. 130: 146. (Cited in U.S. EPA, 1983a)

Rigdon, R.H. and E.G. Rennels. 1964. Effect of feeding benzpyrene on reproduction in the rat. Experientia. 20: 224-226. (Cited in U.S. EPA, 1983b)

Robinson, J.R., J.S. Felton, R.C. Levitt, S.S. Thorgiersson and D.W. Nebert. 1975. Relationship between "aromatic hydrocarbon responsiveness" and the survival times in mice treated with various drugs and environmental compounds. Mol. Pharmacol. 11: 850-865. (Cited in IARC, 1983)

Santodonato, J., P. Howard and D. Basu. 1981. Health and Ecological Assessment of Polynuclear Aromatic Hydrocarbons. Pathotox Publishers, Inc., Park Forest South, IL.

Setala, K. 1954. Experimental chemical carcinogenesis and the influence of solvents. Nature (London). 174: 873. (Cited in Grimmer, 1983)

Smith, J.H., W.R. Mabey, N. Bohonos, et al. 1978. Environmental Pathways of Selected Chemicals in Freshwater Systems. Part II: Laboratory Studies. Environmental Research Laboratory, ORD, U.S. EPA, Athens, GA. EPA 600/7-78-074.

Smyth, H.F., C.P. Carpenter, C.S. Weil, U.C. Pozzani and J.A. Striegel. 1962. Range-finding toxicity data: List VI. Am. Ind. Hyg. Assoc. J. 23: 95-107. (Cited in Santodonato et al., 1981)

Thyssen, J., J. Althoff, G. Kimmerle and U. Mohr. 1981. Inhalation studies with benzo[a]pyrene in Syrian golden hamsters. J. Natl. Cancer Inst. 66(3): 575-577. (Cited in U.S. EPA, 1983a)

U.S. EPA. 1980a. Guidelines and Methodology Used in the Preparation of Health Effects Assessment Chapters of the Consent Decree Water Quality Criteria. Federal Register. 45:79347-79357.

U.S. EPA. 1980b. Ambient Water Quality Criteria for Polynuclear Aromatic Hydrocarbons. Environmental Criteria and Assessment Office, Cincinnati, OH. EPA 440/5-80-069. NTIS PB 81-117806.

U.S. EPA. 1981. Hazard Profile for PAH. Prepared by the Environmental Criteria and Assessment Office, Cincinnati, OH, OHEA, for the Office of Solid Waste and Emergency Response, Washington, DC.

U.S. EPA. 1983a. Review of Toxicologic Data in Support of Evaluation for Carcinogenic Potential of: Benzo[a]pyrene. Prepared by the Carcinogen Assessment Group, OHEA, Washington, DC for the Office of Solid Waste and Emergency Response, Washington, DC.

U.S. EPA. 1983b. Reportable Quantity for Benzo[a]pyrene. Prepared by the Environmental Criteria and Assessment Office, Cincinnati, OH, OHEA for the Office of Solid Waste and Emergency Response, Washington, DC.

U.S. EPA. 1983c. Methodology and Guidelines for Reportable Quantity Determinations Based on Chronic Toxicity Data. Prepared by the Environmental Criteria and Assessment Office, Cincinnati, OH, OHEA for the Office of Solid Waste and Emergency Response, Washington, DC.

Vainio, H., P. Votila, J. Hartiala and O. Pelkonen. 1976. The fate of intratracheally installed benzo[a]pyrene in the isolated perfused rat lung of both control and 20-methylcholanthrene pretreated rats. Res. Comm. Chem. Pathol. Pharmacol. 13: 259271. (Cited in U.S. EPA, 1980b)

Wise, S.A., W.J. Bonnett, F.R. Guenther and W.E. May. 1981. A relationship between reversed-phase C_{18} liquid chromatographic retention and the shape of polycyclic aromatic hydrocarbons. J. Chromatogr. Sci. 19: 457-465.

APPENDIX A

Summary Table for Benzo[a]pyrene

Carcinogenic Potency	Species	Experimental Dose/Exposure	Effect	q_1^* (mg/kg/day) ⁻¹	Reference
Inhalation	hamsters	2.2-9.5 mg/m ³	respiratory tract tumors	6.11	Thyssen et al., 1981
Oral	mice	1-250 ppm	stomach tumors	11.53	Neal and Rigdon, 1967

APPENDIX B

Cancer Data Sheet for Derivation of q_1^*

Compound: Benzo[a]pyrene

Reference: Thyssen et al., 1981

Species, strain, sex: hamsters/Syrian Golden/male

Body weight: 0.12 kg (assumed)

Length of exposure (t_e) = 666.4 days for lower dose and 674.8 days for higher dose and controls

Length of experiment (L_e) = 666.4 days for lower dose and 674.8 days for higher dose and controls

Lifespan of animal (L) = 666.4 days for lower dose and 674.8 days for higher dose and controls

Tumor site and type: respiratory tract/papillomas, papillary polyps and squamous-cell carcinomas

Route, vehicle: inhalation/NaCl vapor in air

Experimental Doses or Exposures*	Input	
	Transformed Dose (mg/kg/day)	Incidence No. Responding/No. Tested (or Examined)
0 mg/m ³	0	0/27
2.2 mg/m ³	0.0892	0/27
9.5 mg/m ³	0.385	9/26

*See following page for conversions

$q_1^*A = 0.731101 \text{ (mg/kg/day)}^{-1}$

$q_1^*H = 6.11 \text{ (mg/kg/day)}^{-1}$

CONVERSIONS

$$0 \text{ mg/m}^3 = 0 \text{ mg/kg/day}$$

$$2.2 \text{ mg/m}^3 \times \left[\frac{10 \text{ weeks}}{95.2 \text{ weeks}} \times \frac{4.5 \text{ hours}}{24 \text{ hours}} \right] + \left(\frac{85.2 \text{ weeks}}{95.2 \text{ weeks}} \times \frac{3 \text{ hours}}{24 \text{ hours}} \right) \times \frac{7 \text{ days}}{7 \text{ days}} \times$$

$$0.037 \text{ m}^3/\text{day} \div 0.12 \text{ kg} \times \frac{666.4 \text{ days}}{666.4 \text{ days}} \times \left(\frac{666.4 \text{ days}}{666.4 \text{ days}} \right)^3 = 0.0892 \text{ mg/kg/day}$$

$$9.5 \text{ mg/m}^3 \times \left[\frac{10 \text{ weeks}}{96.4 \text{ weeks}} \times \frac{4.5 \text{ hours}}{24 \text{ hours}} \right] + \left(\frac{86.4 \text{ weeks}}{96.4 \text{ weeks}} \times \frac{3 \text{ hours}}{24 \text{ hours}} \right) \times \frac{7 \text{ days}}{7 \text{ days}} \times$$

$$0.037 \text{ m}^3/\text{day} \div 0.12 \text{ kg} \times \frac{674.8 \text{ days}}{674.8 \text{ days}} \times \left(\frac{674.8 \text{ days}}{674.8 \text{ days}} \right)^3 = 0.385 \text{ mg/kg/day}$$

DATE

APPENDIX C

Cancer Data Sheet for Derivation of q_1^*

Compound: Benzo[a]pyrene

Reference: Neal and Rigdon, 1967

Species, Strain, Sex: mouse, CFW, M,F

Body weight: 0.034 kg (assumed)

Length of exposure (t_e) = 110 days

Length of experiment (t_e) = 183 days

Lifespan of animal (L) = 630 days

Tumor site and type: stomach, papilloma, carcinomas

Route, vehicle: oral, diet

Experimental Doses or Exposures (ppm)	Transformed Dose (mg/kg/day)*	Incidence
		No. Responding/No. Tested or Examined
0	0	0/289
1	0.078	0/25
10	0.781	0/24
20	1.563	1/23
30	2.344	0/37
40	3.126	1/40
45	3.516	4/40

*Adjusted to reflect treatment on day 110 of a 183-day experimental period.

Unadjusted q_1^* from study = 2.2213189×10^{-2} (mg/kg/day) $^{-1}$

Human q_1^* = 11.53 (mg/kg/day) $^{-1}$

U.S. Environmental Protection Agency
Region V, Library
230 South Dearborn Street
Chicago, Illinois 60604